



BCAD Isoleucine-, Leucine- and Valine-free Powder

1
INFANT
TODDLER

Mead Johnson Nutrition BCAD 1 is to be used only in the dietary management of infants and young children with documented maple syrup urine disease while under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough isoleucine, leucine and valine to support growth using other foods with these amino acids as required. Medical professionals must carefully and constantly supervise use of BCAD 1 with other foods and liquids and adjust the diet based on frequent blood tests.

Instructions for Preparation and Use

Your child's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing product. Powdered products are not sterile and should not be fed to premature infants or persons who might have immune problems unless directed and supervised by your child's medical professional.

Ask your baby's medical professional about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

Your baby's medical professional will provide instructions for the correct amounts of water and powder*. Pour the required amount of water into the bottle. Add the correct amount of powder, cap the bottle and shake vigorously.

*If instructed to use the scoop in the can, each scoop (unpacked and leveled) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, product can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared product if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared product. After feeding begins, use product within one hour or discard.

WARNING: Do not use a microwave oven to warm product. Serious burns may result.

POWDER STORAGE: Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. Use by date on bottom of can.

WARNING: Not for parenteral (I.V.) use.

Mead Johnson
BCAD

DHA & ARA

1
INFANT
TODDLER

INFANT FORMULA & MEDICAL FOOD: Isoleucine-, Leucine- and Valine-free

IRON FORTIFIED POWDER

DHA & ARA

FOR DIETARY MANAGEMENT OF INFANTS AND YOUNG CHILDREN (0-36 MONTHS)
WITH DOCUMENTED MAPLE SYRUP URINE DISEASE

NET WT. 1 LB (454 g) 16 OZ

FROM THE MAKER OF

Enfamil
PRODUCTS

NUTRIENTS	Per 100 g Powder
Calories	500
Protein equivalents, g*	16.2
Fat, g	26
Carbohydrate, g	51
Linoleic acid, mg	4000
Vitamins:	
Vitamin A, IU	1520
Vitamin D, IU	380
Vitamin E, IU	10
Vitamin K, mcg	40
Thiamin (Vitamin B ₁), mcg	1000
Riboflavin (Vitamin B ₂), mcg	1000
Vitamin B ₆ , mcg	1000
Vitamin B ₁₂ , mcg	2
Niacin, mcg	10000
Folic acid (folacin), mcg	100
Pantothenic acid, mcg	3800
Biotin, mcg	38
Vitamin C (ascorbic acid), mg	60
Choline, mg	124
Inositol, mg	86
Minerals:	
Calcium, mg	660
Phosphorous, mg	440
Magnesium, mg	66
Iron, mg	9.6
Zinc, mg	8.6
Manganese, mcg	380
Copper, mcg	860
Iodine, mcg	76
Selenium, mcg	14.1
Sodium, mg	240
Potassium, mg	600
Chloride, mg	500

*WARNING: This protein is incomplete since it does not contain the essential amino acids isoleucine, leucine and valine.

Ingredients: Corn Syrup Solids, Vegetable Oil (Palm Olein, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), AMINO ACIDS (L-glutamine, L-lysine Hydrochloride, Potassium Aspartate, L-proline, L-alanine, L-arginine, L-phenylalanine, L-tyrosine, L-serine, L-threonine, Glycine, L-histidine, L-methionine, L-tryptophan, L-cystine), Modified Corn Starch, Sugar, Calcium Phosphate, and less than 1%: Mortierella Alpina Oil**, Cryptocodinium Cohnii Oil†, Ethyl Vanillin, Sodium Citrate, Potassium Citrate, Magnesium Oxide, Ferrous Sulfate, Potassium Chloride, Zinc Sulfate, Cupric Sulfate, Manganese Sulfate, Sodium Iodide, Sodium Selenite, Choline Chloride, Inositol, Ascorbic Acid, Niacinamide, Calcium Pantothenate, Vitamin B₆ Hydrochloride, Thiamin Hydrochloride, Riboflavin, Vitamin D₃, Folic Acid, Biotin, Vitamin K₁, Vitamin E Acetate, Vitamin A Palmitate, Vitamin B₁₂, Taurine, L-carnitine.

DISTRIBUTED BY:
MEAD JOHNSON & COMPANY, LLC
EVANSVILLE, IN 47721 U.S.A.

©2012 Mead Johnson & Company, LLC

**A Source of Arachidonic Acid (ARA)

†A Source of Docosahexaenoic Acid (DHA)

Filled by weight, not by volume;
some settling may occur.

For additional information, visit our
website at www.enfamil.com

2009154 List No. 892801





OA ISOLEUCINE-, METHIONINE-, THREONINE- & VALINE-FREE POWDER

1
INFANT
TODDLER

Mead Johnson Nutrition OA 1 is to be used only in the dietary management of infants and young children with documented propionic and methylmalonic acidemias while under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough isoleucine, methionine, threonine, and valine to support growth using other foods with these amino acids as required. Medical professionals must carefully and constantly supervise use of OA 1 with other foods and liquids and adjust the diet based on frequent blood tests.

Instructions for Preparation and Use

WARNING: Your child's health depends on carefully following the instructions below. Use only as directed by a medical professional. Improper hygiene, preparation, dilution, use or storage may result in severe harm. Powdered foods are not sterile and should not be fed to premature infants or those persons who might have immune problems unless directed and supervised by your child's medical professional.

Ask your baby's medical professional about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

Your baby's medical professional will provide instructions for the correct amounts of water and powder*. Pour the required amount of water into the bottle. Add the correct amount of powder, cap the bottle and shake vigorously.

*If instructed to use the scoop in the can, each scoop (unpacked and leveled) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, product can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared product if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared product. After feeding begins, use product within one hour or discard.

WARNING: Do not use a microwave oven to warm product. Serious burns may result.

POWDER STORAGE: Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

Use by date on bottom of can.

WARNING: Not for parenteral (I.V.) use.

MeadJohnson®

OA

DHA & ARA

1
INFANT
TODDLER

INFANT FORMULA & MEDICAL FOOD: ISOLEUCINE-, METHIONINE-, THREONINE- & VALINE-FREE

IRON FORTIFIED POWDER

DHA & ARA

FOR DIETARY MANAGEMENT OF INFANTS AND YOUNG CHILDREN (0-36 MONTHS) WITH
DOCUMENTED PROPIONIC AND METHYLMALONIC ACIDEMIAS

FROM THE MAKER OF

Enfamil
PRODUCTS

NET WT. 1 LB (454 g) 16 OZ

NUTRIENTS

	Per 100 g Powder
Calories	500
Protein equivalents, g*	15.7
Fat, g	26
Carbohydrate, g	51
Linoleic acid, mg	4000
Vitamins:	
Vitamin A, IU	1520
Vitamin D, IU	380
Vitamin E, IU	10
Vitamin K, mcg	40
Thiamin (Vitamin B ₁), mcg	1000
Riboflavin (Vitamin B ₂), mcg	1000
Vitamin B ₆ , mcg	1000
Vitamin B ₁₂ , mcg	2
Niacin, mcg	10000
Folic acid (folacin), mcg	100
Pantothenic acid, mcg	3800
Biotin, mcg	38
Vitamin C (ascorbic acid), mg	60
Choline, mg	124
Inositol, mg	86
Minerals:	
Calcium, mg	660
Phosphorus, mg	440
Magnesium, mg	66
Iron, mg	9.6
Zinc, mg	8.6
Manganese, mcg	380
Copper, mcg	860
Iodine, mcg	76
Selenium, mcg	14.1
Sodium, mg	240
Potassium, mg	560
Chloride, mg	480

*WARNING: This protein is incomplete since it does not contain the essential amino acids isoleucine, methionine, threonine, and valine.

Ingredients: Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), AMINO ACIDS (L-alanine, L-leucine, L-lysine Hydrochloride, Potassium Aspartate, L-proline, L-arginine, Glycine, L-phenylalanine, L-tyrosine, L-serine, L-histidine, L-tryptophan, L-cystine), Modified Corn Starch, Sugar, Calcium Phosphate, and less than 1%: Mortierella Alpina Oil*, Cryptocodium Cohnii Oil*, Ethyl Vanillin, Sodium Citrate, Potassium Citrate, Magnesium Oxide, Ferrous Sulfate, Potassium Chloride, Zinc Sulfate, Cupric Sulfate, Manganese Sulfate, Sodium Iodide, Sodium Selenite, Choline Chloride, Inositol, Ascorbic Acid, Niacinamide, Calcium Pantothenate, Vitamin B₆ Hydrochloride, Thiamin Hydrochloride, Riboflavin, Vitamin D₃, Folic Acid, Biotin, Vitamin K₁, Vitamin E Acetate, Vitamin A Palmitate, Vitamin B₁₂, Taurine, L-carnitine.

DISTRIBUTED BY:

MEAD JOHNSON & COMPANY, LLC
EVANSVILLE, IN 47721 U.S.A.
©2012 Mead Johnson & Company, LLC

**A Source of Arachidonic Acid (ARA)


*A Source of Docosahexaenoic Acid (DHA)

Filled by weight, not by volume;
some settling may occur.

For additional information, visit our
website at www.enfamil.com

2009175 List No. 893201





PFD PROTEIN- AND AMINO ACID-FREE POWDER

TODDLER

Mead Johnson Nutrition PFD TODDLER is to be used only in the dietary management of young children with documented amino acid metabolic disorders while under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough protein and amino acids to support growth using other foods with protein as required. Medical professionals must carefully and constantly supervise use of PFD TODDLER with other foods and liquids and adjust the diet based on frequent blood tests.

Instructions for Preparation and Use

Your medical professional will provide the correct amount of powder to mix with water for consumption.*

It is important to follow the directions below.

Measure the correct amount of water into a suitable container for mixing. Then add the required amount of PFD TODDLER. Mix well until blended. Consume the prepared beverage immediately or cover and refrigerate. Use within 48 hours of preparation. Mix before drinking.

* If instructed to use the scoop in the can, each unpacked, level scoop delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

POWDER STORAGE: Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

Use by date on can end.

WARNING: Not for parenteral (I.V.) use.

MeadJohnson®

PFD

DHA & ARA

MEDICAL FOOD: PROTEIN- AND AMINO ACID-FREE IRON FORTIFIED POWDER

DHA & ARA

FOR DIETARY MANAGEMENT OF YOUNG CHILDREN (1-3 YEARS)
WITH DOCUMENTED INBORN ERRORS OF
AMINO ACID METABOLISM

TODDLER

FROM THE MAKER OF

Enfamil
PRODUCTS™

NET WT. 1 LB (454 g) 16 OZ

NUTRIENTS

	Per 100 g Powder
Calories	530
Protein equivalents, g*	0
Fat, g	32
Carbohydrate, g	60
Linoleic acid, mg	4900
Vitamins:	
Vitamin A, IU	1840
Vitamin D, IU	450
Vitamin E, IU	13.2
Vitamin K, mcg	53
Thiamin (Vitamin B ₁), mcg	1320
Riboflavin (Vitamin B ₂), mcg	1210
Vitamin B ₆ , mcg	1210
Vitamin B ₁₂ , mcg	2.6
Niacin, mcg	12100
Folic acid (folacin), mcg	132
Pantothenic acid, mcg	4600
Biotin, mcg	47
Vitamin C (ascorbic acid), mg	74
Choline, mg	135
Inositol, mg	109
Minerals:	
Calcium, mg	790
Phosphorous, mg	530
Magnesium, mg	79
Iron, mg	10.5
Zinc, mg	10.5
Manganese, mcg	470
Copper, mcg	1050
Iodine, mcg	90
Selenium, mcg	15.8
Sodium, mg	280
Potassium, mg	690
Chloride, mg	530

*WARNING: This product is nutritionally incomplete since it does not contain any protein or amino acids.

Ingredients: Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), Modified Corn Starch, Sugar, Calcium Phosphate, and less than 1%: Mortierella Alpina Oil**, Cryptocodinium Cohnii Oil*, Ethyl Vanillin, Sodium Citrate, Potassium Chloride, Potassium Citrate, Magnesium Oxide, Ferrous Sulfate, Zinc Sulfate, Cupric Sulfate, Manganese Sulfate, Sodium Iodide, Sodium Selenite, Choline Chloride, Inositol, Ascorbic Acid, Niacinamide, Calcium Pantothenate, Vitamin B₆ Hydrochloride, Thiamin Hydrochloride, Riboflavin, Vitamin D₃, Folic Acid, Biotin, Vitamin K₁, Vitamin E Acetate, Vitamin A Palmitate, Vitamin B₁₂, Taurine, L-carnitine.

DISTRIBUTED BY:
MEAD JOHNSON & COMPANY, LLC
EVANSVILLE, IN 47721 U.S.A.
©2012 Mead Johnson & Company, LLC

**A Source of Arachidonic Acid (ARA)
*A Source of Docosahexaenoic Acid (DHA)

Filled by weight, not by volume;
some settling may occur.

For additional information, visit our
website at www.enfamil.com

2009173

List No. 892701



Formula for Special Medical Purpose
TO BE TAKEN UNDER MEDICAL ADVICE



Imported by:
Mead Johnson Nutrition (India) Pvt. Ltd.
3rd Floor, Piramal Tower,
Peninsula Corporate Park,
Lower Parel, Mumbai - 400013

Toll-free No.: 1800220118
Email: meadjohnsonnuture.india@mjn.com

Mfd. by:
Mead Johnson & Company, LLC
Evansville, IN 47721 USA

Product of the U.S.A.

Lic.No. 10012022000280

Diet 4 Life
An FSSAI Initiative

Mfg. Date
MRP in Rs.
Inclusive of all taxes

Expiry Date/Batch No. see bottom
This is not an infant milk substitute or infant food

PHENYL-FREE[®] PHENYLALANINE-FREE POWDER

1
INFANT
TODDLER

Mead Johnson Nutrition PHENYL-FREE 1 is to be used only in the dietary management of infants and young children with documented phenylketonuria while under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough phenylalanine to support growth using other foods with this amino acid as required. Medical professionals must carefully and constantly supervise use of PHENYL-FREE 1 with other foods and liquids and adjust the diet based on frequent blood tests.

Instructions for Preparation and Use

Your child's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing product. Powdered products are not sterile and should not be fed to premature infants or persons who might have immune problems unless directed and supervised by your child's medical professional.

Ask your baby's medical professional about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

Your baby's medical professional will provide instructions for the correct amounts of water and powder*. Pour the required amount of water into the bottle. Add the correct amount of powder, cap the bottle and shake vigorously.

*If instructed to use the scoop in the can, each scoop (unpacked and leveled) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, product can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared product if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared product. After feeding begins, use product within one hour or discard.

WARNING: Do not use a microwave oven to warm product. Serious burns may result.

POWDER STORAGE: Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

Use by date on bottom of can.

WARNING: Not for parenteral (I.V.) use.

MeadJohnson[®]

DHA & ARA

PHENYL-FREE[®] INFANT FORMULA & MEDICAL FOOD: PHENYLALANINE-FREE IRON FORTIFIED POWDER

1
INFANT
TODDLER

DHA & ARA

FOR DIETARY MANAGEMENT OF INFANTS AND YOUNG CHILDREN (0-36 MONTHS) WITH DOCUMENTED PHENYLKETONURIA

FROM THE MAKER OF

Enfamil[®] PRODUCTS

NET WT. 1 LB (454 g) 16 OZ

NUTRIENTS	Per 100 g Powder
Calories	500
Protein equivalents, g*	16.2
Fat, g	26
Carbohydrate, g	51
Linoleic acid, mg	4000
Vitamins:	
Vitamin A, IU	1520
Vitamin D, IU	380
Vitamin E, IU	10
Vitamin K, mcg	40
Thiamin (Vitamin B ₁), mcg	1000
Riboflavin (Vitamin B ₂), mcg	1000
Vitamin B ₆ , mcg	1000
Vitamin B ₁₂ , mcg	2
Niacin, mcg	10000
Folic acid (folacin), mcg	100
Pantothenic acid, mcg	3800
Biotin, mcg	38
Vitamin C (ascorbic acid), mg	60
Choline, mg	124
Inositol, mg	86
Minerals:	
Calcium, mg	660
Phosphorus, mg	440
Magnesium, mg	66
Iron, mg	9.6
Zinc, mg	8.6
Manganese, mcg	380
Copper, mcg	860
Iodine, mcg	76
Selenium, mcg	14.1
Sodium, mg	240
Potassium, mg	560
Chloride, mg	430

*WARNING: This protein is incomplete since it does not contain the essential amino acid phenylalanine.

Ingredients: Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), Amino Acids (L-glutamine, L-leucine, Potassium Aspartate, L-lysine Hydrochloride, L-tyrosine, L-proline, L-valine, L-isoleucine, L-alanine, L-arginine, L-threonine, L-serine, Glycine, L-histidine, L-methionine, L-tryptophan, L-cystine), Modified Corn Starch, Sugar, Calcium Phosphate, and less than 1%: Mortierella Alpina Oil**, Cryptocodinium Cohnii Oil†, Ethyl Vanillin, Sodium Citrate, Potassium Citrate, Magnesium Oxide, Ferrous Sulfate, Potassium Chloride, Zinc Sulfate, Cupric Sulfate, Manganese Sulfate, Sodium Iodide, Sodium Selenite, Choline Chloride, Inositol, Ascorbic Acid, Niacinamide, Calcium Pantothenate, Vitamin B₆ Hydrochloride, Thiamin Hydrochloride, Riboflavin, Vitamin D₃, Folic Acid, Biotin, Vitamin K₁, Vitamin E Acetate, Vitamin A Palmitate, Vitamin B₁₂, Taurine, L-carnitine.

DISTRIBUTED BY:
MEAD JOHNSON & COMPANY, LLC
EVANSVILLE, IN 47721 U.S.A.
©2012 Mead Johnson & Company, LLC

**A Source of Arachidonic Acid (ARA)
†A Source of Docosahexaenoic Acid (DHA)

Filled by weight, not by volume;
some settling may occur.

For additional information, visit our
website at www.enfamil.com

2009171 List No. 892601





TYROS PHENYLALANINE- AND TYROSINE-FREE POWDER

1
INFANT
TODDLER

Mead Johnson Nutrition TYROS 1 is to be used only in the dietary management of infants and young children with documented tyrosinemia while under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough phenylalanine and tyrosine to support growth using other foods with these amino acids as required. Medical professionals must carefully and constantly supervise use of TYROS 1 with other foods and liquids and adjust the diet based on frequent blood tests.

Instructions for Preparation and Use

Your child's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing product. Powdered products are not sterile and should not be fed to premature infants or persons who might have immune problems unless directed and supervised by your child's medical professional.

Ask your baby's medical professional about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

Your baby's medical professional will provide instructions for the correct amounts of water and powder*. Pour the required amount of water into the bottle. Add the correct amount of powder, cap the bottle and shake vigorously.

*If instructed to use the scoop in the can, each scoop (unpacked and leveled) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, product can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared product if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared product. After feeding begins, use product within one hour or discard.

WARNING: Do not use a microwave oven to warm product. Serious burns may result.

POWDER STORAGE: Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

Use by date on bottom of can.

WARNING: Not for parenteral (I.V.) use.

MeadJohnson

TYROS

DHA & ARA

1
INFANT
TODDLER

INFANT FORMULA & MEDICAL FOOD: PHENYLALANINE- AND TYROSINE-FREE IRON FORTIFIED POWDER

DHA & ARA

FOR DIETARY MANAGEMENT OF INFANTS AND YOUNG CHILDREN (0-36 MONTHS) WITH DOCUMENTED TYROSINEMIA

FROM THE MAKER OF

Enfamil
PRODUCTS

NET WT. 1 LB (454 g) 16 OZ

NUTRIENTS	Per 100 g Powder
Calories	500
Protein equivalents, g*	16.7
Fat, g	26
Carbohydrate, g	51
Linoleic acid, mg	4000
Vitamins:	
Vitamin A, IU	1520
Vitamin D, IU	380
Vitamin E, IU	10
Vitamin K, mcg	40
Thiamin (Vitamin B ₁), mcg	1000
Riboflavin (Vitamin B ₂), mcg	1000
Vitamin B ₆ , mcg	1000
Vitamin B ₁₂ , mcg	2
Niacin, mcg	10000
Folic acid (folacin), mcg	100
Pantothenic acid, mcg	3800
Biotin, mcg	38
Vitamin C (ascorbic acid), mg	60
Choline, mg	124
Inositol, mg	86
Minerals:	
Calcium, mg	660
Phosphorous, mg	440
Magnesium, mg	66
Iron, mg	9.6
Zinc, mg	8.6
Manganese, mcg	380
Copper, mcg	860
Iodine, mcg	76
Selenium, mcg	14.1
Sodium, mg	240
Potassium, mg	610
Chloride, mg	430

*WARNING: This protein is incomplete since it does not contain the essential amino acids phenylalanine and tyrosine.

Ingredients: Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), AMINO ACIDS (L-glutamine, L-leucine, Potassium Aspartate, L-lysine Hydrochloride, L-proline, L-valine, L-alanine, L-isoleucine, L-arginine, L-threonine, L-serine, Glycine, L-histidine, L-methionine, L-tryptophan, L-cystine), Modified Corn Starch, Sugar, Calcium Phosphate, and less than 1%: Mortierella Alpina Oil**, Cryptocodinium Cohnii Oil*, Ethyl Vanillin, Sodium Citrate, Potassium Citrate, Magnesium Oxide, Ferrous Sulfate, Potassium Chloride, Zinc Sulfate, Cupric Sulfate, Manganese Sulfate, Sodium Iodide, Sodium Selenite, Choline Chloride, Inositol, Ascorbic Acid, Niacinamide, Calcium Pantothenate, Vitamin B₆ Hydrochloride, Thiamin Hydrochloride, Riboflavin, Vitamin D₃, Folic Acid, Biotin, Vitamin K₁, Vitamin E Acetate, Vitamin A Palmitate, Vitamin B₁₂, Taurine, L-carnitine.

DISTRIBUTED BY:
MEAD JOHNSON & COMPANY, LLC
EVANSVILLE, IN 47721 U.S.A.
©2012 Mead Johnson & Company, LLC

**A Source of Arachidonic Acid (ARA)
*A Source of Docosahexaenoic Acid (DHA)

Filled by weight, not by volume;
some settling may occur.

For additional information, visit our
website at www.enfamil.com

2009172 List No. 893001



Formula for Special Medical Purpose

TO BE TAKEN UNDER MEDICAL ADVICE

Imported by:
Mead Johnson Nutrition (India) Pvt. Ltd.
3rd Floor, Pinnacle Tower,
Peninsula Corporate Park,
Lower Parel, Mumbai - 400013

Toll-free No.: 1800220118
Email: meadjohnsonnutrition.india@mjn.com

Mfg. by:
Mead Johnson & Company, LLC
Evansville, IN 47721 USA

Product of the U.S.A.

Lic. No. 10012022000280



fssai



Diet 4Life

An FSSAI Initiative

Mfg. Date
MRP in Rs.
Inclusive of all taxes

Expiry Date/Batch No. see bottom

This is not an Infant milk substitute of any kind



PURAMINO™ is a hypoallergenic, amino acid-based formula for infants and toddlers with severe cow's milk protein and multiple food allergies. It is nutritionally complete for 0-12 months; dietary reference intakes include a recommendation for fiber >12 months of age. It is also suitable for conditions requiring an elemental diet such as protein maldigestion, malabsorption/chronic diarrhea, short bowel syndrome and eosinophilic esophagitis.

If you have a question, we are here for you. Call us toll free: 1-800-BABY123,
8 am to 4:30 pm Monday-Saturday, Central Time. Or visit Entamil.com

INSTRUCTIONS FOR PREPARATION & USE

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

1. Wash hands thoroughly with soap and water before preparing formula.



2. Pour desired amount of water into the bottle. Add powder.



3. Cap bottle and SHAKE WELL.



Use scoop in can to measure powder. Store DRY scoop in this can.

For every 1 fl oz of water, add 1 unpacked level scoop of powder (4.5 g). Each scoop adds about 0.1 fl oz to the amount of prepared formula. For example, adding 3 unpacked level scoops of powder to 3 fl oz of water will make about 3.3 fl oz of formula.

WARNING: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in

refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula.

After feeding begins, use formula within one hour or discard.

Powder Storage: Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

USE BY DATE ON BOTTOM OF CAN

CORTE ADU PARALEL LAS INSTRUCCIONES EN ESPAÑOL

PURAMINO™ DHA & ARA

Hypoallergenic Infant Formula amino acid-based powder with iron

For infants and toddlers with severe
cow's milk protein and multiple food allergies

Also suitable for conditions requiring an elemental diet such as protein maldigestion,
malabsorption/chronic diarrhea, short bowel syndrome and eosinophilic esophagitis

0-24 months

NET WT 14.1 OZ (400 g)

From the
maker of

Nutramigen

U D M

NUTRIENTS (Normal Dilution)	Per 100 Calories (5 fl oz)	
Protein Equivalent	g	2.8
Fat	g	5.3
Carbohydrate	g	10.6
Water	g	133
Linoleic acid	mg	860
Vitamins:		
A	IU	300
D	IU	50
E	IU	2
K	mcg	9
Thiamin (Vitamin B ₁)	mcg	80
Riboflavin (Vitamin B ₂)	mcg	90
B ₆	mcg	60
B ₁₂	mcg	0.3
Niacin	mcg	1000
Folic acid (Folacin)	mcg	16
Pantothenic acid	mcg	500
Biotin	mcg	3
C (Ascorbic acid)	mg	12
Choline	mg	24
Inositol	mg	17
Minerals:		
Calcium	mg	94
Phosphorus	mg	52
Magnesium	mg	11
Iron	mg	1.8
Zinc	mg	1
Manganese	mcg	25
Copper	mcg	75
Iodine	mcg	15
Selenium	mcg	2.8
Sodium	mg	47
Potassium	mg	110
Chloride	mg	86

INGREDIENTS: CORN SYRUP SOLIDS (48%), AMINO ACIDS (L-ASPARTIC ACID, L-LEUCINE, L-PROLINE, L-LYSINE, HYDROCHLORIDE, L-ALANINE, MONOSODIUM GLUTAMATE, L-VALINE, L-ISOLEUCINE, L-THREONINE, L-SERINE, L-TYROSINE, L-ARGININE, L-PHENYLALANINE, GLYCINE, L-HISTIDINE, L-CYSTINE, L-TRYPTOPHAN, L-METHIONINE) (18%), HIGH OLEIC SUNFLOWER OIL (10%), MEDIUM CHAIN TRIGLYCERIDES (MCT) OIL (9%), SOY OIL (8%), MODIFIED TAPIOCA STARCH (3%), AND LESS THAN 2% MORTIERELLA ALPINA OIL¹, CRYPTOCODINIUM COHNI OIL¹, CHOLINE CHLORIDE, INOSITOL, ASCORBIC ACID, NIACINAMIDE, CALCIUM PANTOTHENATE, RIBOFLAVIN, THIAMIN HYDROCHLORIDE, VITAMIN B₆ HYDROCHLORIDE, VITAMIN D₃, FOLIC ACID, VITAMIN K₁, BIOTIN, TAURINE, VITAMIN E ACETATE, L-CARNITINE, POTASSIUM CITRATE, CALCIUM CITRATE, SODIUM CITRATE, POTASSIUM CHLORIDE, CALCIUM PHOSPHATE, MAGNESIUM PHOSPHATE, FERRIC SULFATE, ZINC SULFATE, COPPER SULFATE, MANGANESE SULFATE, SODIUM IODIDE, SODIUM SELENITE.

MEAD JOHNSON & COMPANY, LLC
EVANSVILLE, IN 47721 U.S.A.

MeadJohnson
Nutrition

© 2016 Mead Johnson & Company, LLC

¹A SOURCE OF ARACHIDONIC ACID (ARA)

¹A SOURCE OF DOCOSAHEXAENOIC ACID (DHA)

Filled by weight, not by volume; some settling may occur.

Makes 98 fl oz

Patent Pending

2036725



Nutramigen LGG* is a hypoallergenic formula designed to meet all the daily needs of babies who are allergic to the intact proteins in cow's milk and soy formulas, as well as other foods. Nutramigen LGG is also lactose-free.

Clinically proven, fast management of cow's milk allergies

- The proteins in Nutramigen LGG are extensively hydrolyzed into tiny pieces that they are unlikely to trigger an allergic reaction.
- Contains Lactobacillus GG (LGG), a combination^a that's clinically proven to improve the dietary management of cow's milk allergy and helps your baby return to a normal diet faster.
- Nutramigen, a brand that has been clinically proven safe and effective in managing cow's milk protein allergy for more than 70 years and in over 70 clinical trials.


^a The combination of extensively hydrolyzed protein and LGG

WARNING/CAUTION

- Careful & hygienic preparation of infant food / infant milk substitute is most essential for health. Do not use fewer scoops than directed since diluted feeding will not provide adequate nutrients needed by your infant. Do not use more scoops than directed since concentrated feed will not provide the water needed by your infant.
- Infant milk substitute should be used only on the advice of a health worker as to the need for its use and the proper method of its use.
- Infant milk substitute is not the sole source of nourishment of an infant. Extensively hydrolyzed formula tastes and smells different from routine formula. Clinical studies show that most babies adapt Nutramigen quickly and do not have a problem with the taste. Please allow your baby to get used to it.

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby. Ask your baby's doctor about the need to use cooled, boiled water for mixing and the need to boil clean utensils, bottles and nipples in water before use.

INSTRUCTIONS FOR PREPARATION & USE



MFG DATE/BATCH NO/EXPIRY DATE/MRP IN RS.
INCLUSIVE OF ALL TAXES, SEE BOTTOM

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in can to measure powder. Store DRY scoop in this can.

Age of Infant (Weeks)	Number of feedings per day	Level Scoops	For single feeding Boiled Water (40°C)	For single feeding Boiled Water (40°C)
0-2 (Weeks)	6	3	90	3
2-4 (Weeks)	7	3	90	3
4-8 (Weeks)	8	3	90	3
8-12 (Weeks)	5	5	150	5
12-24 (Weeks)	5	5	150	5
6 months onward	3-5	6	180	6
	2-4	8	240	8

One unpacked level scoop (4.5 g) of Nutramigen LGG powder for each 30 ml (1 fl. oz.) of water.

Powder Storage: Store in a cool & dry place. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

Important Notice:
Do not use if the aluminium lid is open or missing.

fsai

Lic.No. 10012022000280

TO BE TAKEN UNDER MEDICAL ADVICE



Hypoallergenic Formula

For the dietary management of cow's milk protein allergy

- Colic and eczema due to cow's milk allergy
- Soy allergy
- Lactose-free
- Extensively-hydrolyzed protein

Net weight: 400g

Formula for Special Medical Purpose for Infants Proprietary Food

IMPORTANT NOTICE: MOTHER'S MILK IS BEST FOR YOUR BABY



INGREDIENTS: CORN SYRUP SOLIDS, EDIBLE VEGETABLE OILS (PALM OLEIN, COCONUT, SOYBEAN AND HIGH OLEIC SUNFLOWER), EXTENSIVELY-HYDROLYZED CASEIN, MODIFIED CORN STARCH, MINERALS, ARACHIDONIC ACID (ARA) FROM MORTIERELLA ALPINA, DOCOSAHEXAENOIC ACID (DHA) from CRYPTOCODINIUM COHNII, VITAMINS, L-CYSTINE, INOSITOL, L-TYROSINE, L-TRYPTOPHAN, TAURINE, LACTOBACILLUS RHAMNOSUS GG (LGG), L-CARNITINE AND ASCORBIC ACID.

WARNING: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard. **Caution:** Do not warm the prepared formula. Warming may limit the benefits of LGG. Infants over six months of age should receive supplemental foods in addition to the formula. As your baby grows, consult your doctor regarding the addition of complementary foods to your baby's diet. After consultation with doctor, baby can continue with Nutramigen LGG from 6 months onwards as part of a diversified diet.

Imported by: Mead Johnson Nutrition (India) Pvt. Ltd., 3rd Floor, Piramal Tower, Peninsula Corporate Park, Ganpatrao Kadam Marg, Lower Parel, Mumbai - 400013

Mfd. by: Mead Johnson B. V., Middenkampung 2, P.O. Box 40003, 6504 AA Nijmegen, The Netherlands

Write to us at: MJN Customer Services, P.O. Bag No. 16303, Mumbai - 400013
E-mail us at: meadjohnsonnurture.india@mjn.com
Call us at toll-free no.: 1800 220 118

Product of the Netherlands

© 2017 Mead Johnson & Company


*LGG is a registered

trademark of Valio Ltd (Finland)




Nutrients	Per 100 g	Per 100 kcal
Food energy, kcal	500	100
Protein, g	14	2.8
Total Fat, g	25	5
Saturated Fatty Acids, g	10.7	2.1
Monounsaturated Fatty Acids, g	9.2	1.84
Polyunsaturated Fatty Acids, g	4.9	0.98
Transfatty Acids, mg	0.0	0.0
Linoleic Acid, mg	4500	900
α-Linolenic Acid, mg	340	68
Arachidonic Acid (ARA), mg	170	34
Docosahexaenoic Acid (DHA), mg	85	17
Carbohydrates, g	55	11
of which sugar (sucrose), g	0.0	0.0
Minerals		
Calcium, mg	570	113
Phosphorus, mg	390	78
Iodine, mcg	105	21
Iron, mg	9	1.8
Magnesium, mg	50	10
Zinc, mg	3.5	0.7
Copper, mcg	380	75
Manganese, mcg	300	60
Sodium, mg	240	47
Potassium, mg	610	122
Chloride, mg	480	96
Selenium, mcg	11	2.2
Vitamins		
Vitamin A, mcg RE	450	90
Vitamin D, mcg	7.5	1.5
Vitamin E, mg α-TE	6.7	1.34
Vitamin C, mg	105	21
Folic Acid, mcg	80	16
Thiamin, mcg	400	80
Riboflavin, mcg	450	90
Niacin, mcg	5000	1000
Vitamin B6, mcg	300	60
Vitamin B12, mcg	1.5	0.3
Biotin, mcg	15	3
Pantothenic Acid, mcg	2500	500
Vitamin K, mcg	65	13
Choline, mg	120	24
Inositol, mg	85	17
Taurine, mg	30	6
L-Carnitine, mg	12.5	2.5
Molybdenum, mcg	28	5.6
Chromium, mcg	11	2.2
Probiotic (LGG), cfu	1x10 ⁸	2x10 ⁷

Formula for Special Medical Purpose
TO BE TAKEN UNDER MEDICAL ADVICE

 **Imported by:**
Mead Johnson Nutrition (India) Pvt. Ltd.
3rd Floor, Piramal Tower,
Peninsula Corporate Park,
Lower Parel, Mumbai - 400013

Lic.No. 10012022000280 **Product of the Netherlands**

Toll-free No.: 1800220118
Email: meadjohnsonnuture.india@mjn.com
Batch No./Mfg. Date/Exp. Date, see bottom

 **An FSSAI Initiative**

MRP in Rs.
Inclusive of all taxes

This is not an Infant milk substitute or Infant food

Pregestimil Powder – Food for Special Medical Purposes

Pregestimil is especially designed for infants with allergy and/or fat maldigestion/ malabsorption with symptoms such as protracted diarrhoea, gastrointestinal symptoms or failure to thrive. It is a **nutritionally complete** formula with extensively hydrolysed protein and can be used as a sole source of nourishment. After 6 months, your baby can continue with Pregestimil as part of a mixed diet. Pregestimil contains a high proportion of Medium Chain Triglycerides, which are easily absorbed. Pregestimil is lactose free and was the first formula for allergic babies with the recommended DHA (Omega 3) level to support normal visual development*. DHA is also present in breast milk. Extensively hydrolysed formula tastes and smells different from standard formula, so **allow your baby to get used to it**.

* The beneficial effect is obtained with a daily intake of 100mg of DHA.

For more information visit: www.nutramigen.co.uk Or call Careline: +44 (0) 1895 230575

Feeding guidelines, unless your healthcare professional advises you otherwise.
Use the chart below for the correct amounts of boiled water and powder

Age	Approx. weight kg / lbs	Number of bottles/24h	Water/feed (ml/fl.oz)	+ scoops *
0 - 2 weeks	< 3.5 kg / 6½ - 7½	6	90 / 3	3
2 - 4 weeks	3.5 - 4 kg / 7½ - 8½	5 - 6	120 / 4	4
1 - 2 months	4 - 5 kg / 8½ - 11	5	150 / 5	5
2 - 4 months	5 - 6.5 kg / 11 - 14	4 - 5	180 / 6	6
> 4 months	> 6.5 kg / >14	4	210 / 7	7

* Only use scoop in can. Do not press powder into scoop, level off with the back of a clean, dry knife. 1 level scoop (4.5 g) to each 30 ml of water.

Instructions for preparation and use:

1. Wash hands thoroughly. Clean bottles, teats, caps and utensils in water (1 minute at a rolling boil).
2. Boil fresh water. Leave to cool for 30 minutes prior to mixing. Do not use softened water. Pour desired amount of water into clean bottle.
3. Add required number of level scoops of powder (see feeding guidelines). Store dry scoop in the can.
4. Cap bottle, roll and shake well until powder dissolves completely. Check temperature on the inside of your wrist.
5. Once prepared, formula can spoil quickly. Feed immediately after preparation or cover and store in a refrigerator at 2-4°C for no longer than a total of 24 hours. Do not freeze prepared formula and do not use if unrefrigerated for more than a total of 2 hours. Use formula that has been in contact with baby's mouth within 1 hour and do not refrigerate. Throw away prepared formula left in bottle and clean utensils. Do not heat feeds in a microwave.

Mead Johnson Nutrition, 271 High Street, Uxbridge, UB8 1LQ
Produced by Mead Johnson Nutrition, Middenkampweg 2,
6545 CJ Nijmegen, The Netherlands
*Trademark of Mead Johnson & Company
©2014 Mead Johnson & Company. All rights reserved.

Best before:
see bottom of can.
Packaged in a protective
atmosphere.



Pregestimil^{*}

LIPIL

Hypoallergenic Formula with MCT oil

For the Dietary Management of Cow's Milk or Soya Protein Allergy

Protracted Diarrhoea and Fat Malabsorption

400g e

MeadJohnson
Nutrition

From birth onwards

Pregestimil

Important notice: Breast milk is the best nutrition for babies. The decision to discontinue breastfeeding may be difficult to reverse and the introduction of partial bottle-feeding may reduce breast milk supply. Pregestimil is intended for the dietary management of cow's milk/soya protein allergy and/or fat malabsorption with symptoms such as protracted diarrhoea and must be used under medical supervision. Suitable for babies from birth onwards when bottle fed, it can be used as a sole source of nourishment, or as part of a mixed diet after 6 months. Babies fed with this formula may produce frequent loose stools of a greenish colour. **Warning:** For your baby's health we advise you to carefully follow the instructions for preparation and use. Proper hygiene, preparation, dilution, use and storage are important when preparing feeding bottles. Powdered formulas intended for infants are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Always ask your baby's healthcare professional which formula is appropriate for your baby. **Failure to follow these instructions could result in severe harm. For enteral use only. Powder storage:** Store can in a cool dry place. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. **Ingredients:** Glucose syrup, vegetable oils (coconut medium chain triglycerides (15.2%), soya, high oleic sunflower), extensively hydrolysed casein (from milk), modified corn starch, calcium phosphate, calcium citrate, potassium citrate, potassium chloride, single cell oils (ARA from Mortierella alpina and DHA from Cryptocodinium cohnii), L-cystine, L-tyrosine, sodium L-ascorbate, choline chloride, sodium citrate, inositol, magnesium oxide, L-tryptophan, ferrous sulphate, taurine, DL-alpha tocopheryl acetate, L-carnitine, zinc sulphate, ascorbic acid, nicotinamide, calcium pantothenate, cupric sulphate, retinyl palmitate, manganese sulphate, thiamin hydrochloride, riboflavin, cholecalciferol, pyridoxine hydrochloride, sodium iodide, folic acid, sodium molybdate, chromic chloride, phytomenadione, D-biotin, sodium selenite, cyanocobalamin.
May not be suitable for some cultural groups

Nutrient	Unit	Per 100 g powder	Per 100 ml reconstituted
Energy	kJ (kcal)	2100 (500)	280 (68)
Fat of which:	g	28	3.8
Saturates	g	16.8	2.3
MCT	g	15.2	2.1
Linoleic acid	g	5.6	0.76
α-Linolenic acid	mg	700	95
Carbohydrate, of which	g	51	6.9
Sugars	g	6.2	0.84
Protein	g	14	1.89
Salt	g	0.55	0.075
Vitamin A	µg RE	570	77
Vitamin D	µg	9.3	1.25
Vitamin E	mg α-TE	13.4	1.81
Vitamin K	µg	60	8.1
Vitamin C	mg	96	13
Thiamin	mg	0.4	0.054
Riboflavin	mg	0.45	0.061
Niacin	mg NE	5	0.68
Vitamin B6	mg	0.3	0.041
Folic Acid	µg	80	10.8
Vitamin B12	µg	1.5	0.2
Biotin	µg	15	2
Pantothenic Acid	mg	2.5	0.34
Sodium	mg (mmol)	220 (9.3)	29 (1.26)
Potassium	mg (mmol)	550 (14.1)	74 (1.9)
Chloride	mg (mmol)	430 (12.1)	58 (1.64)
Calcium	mg (mmol)	580 (14.3)	78 (1.94)
Phosphorus (PO ₄)	mg (mmol)	380 (12.1)	51 (1.64)
Magnesium	mg (mmol)	57 (2.3)	7.5 (0.31)
Iron	mg	9	1.22
Zinc	mg	5	0.68
Copper	mg	0.38	0.051
Manganese	mg	0.3	0.041
Selenium	µg	11	1.49
Chromium	µg	11	1.49
Molybdenum	µg	28	3.8
Iodine	µg	105	14.2
Osmolality	251 mOsm/L		
Arachidonic acid (ARA)	mg	170	23
Docosahexaenoic acid (DHA)	mg	85	11.5
Choline	mg	121	16
Inositol	mg	65	11.5
L-Carnitine	mg	12.5	1.69
Taurine	mg	30	4.1